Copper Ultra

Contents: 5 Litres e

Contains: Copper 250 g/L (as the Oxychloride)

An Inorganic Micronutrient for foliar application to prevent and correct Copper deficiency

Rates of Use:

Deficiency Status
Maintenance
Moderate
Severe

D.50 litres/ha
0.50 litres/ha
0.50 litres/ha & repeat
as necessary

PRECAUTIONS:

Harmful by inhalation and if swallowed. May cause sensitisation by skin contact. Very toxic to aquatic organisms, may cause long-term adverse effects in the aquatic environment. Do not breathe spray. Avoid contact with skin. Wear suitable protective clothing, gloves and eye / face protection. Use appropriate container to avoid environmental contamination. This material and its container must be disposed of as hazardous waste. Avoid release to the environment. Refer to special instructions / safety data sheets. Contains 1,2-benzisothiazolin-3-one. May produce an alleroic reaction.

CONDITIONS OF SUPPLY: All goods supplied by us are of high grade and we believe them to be suitable, but as we cannot we services control over their storage, mixing or use, all conditions and warranties, statutory or otherwise as to the quality or fitness for any purpose of our goods are excluded and no responsibility will be accepted by us for any damage or injury whatsoever arisins from their storage. handling, accidication or use.

Copper OXY 500 is a Trademark of De Sangosse Ltd.

Crop Recommendations and Growth Stages

Cereals 2 leaf stage to 2nd node

Maize 4-10 true leaves Oilseed Rape 4-8 true leaves

Top Fruit At petal fall
Vegetables When leaf area is sufficient

When leaf area is sufficient



Others

HARMFUL



DANGEROUS FOR THE ENVIRONMENT

Water Volumes:

Apply from 200 - 300 litres/ha

TANK MIXING:

Premixing is not necessary. Maintain agitation throughout the spraying period and DO NOT allow the mix to stand for long periods before use. Add Copper Oxy 500 to the spray tank as the last component. Spraying equipment should be thoroughly cleaned and washed out with clean water after completion of spraying.



SHAKE WELL BEFORE USE